arXiv:0808.1847v1 [hep-ph] 13 Aug 2008

Parton Distributions and QCD at LHCb

R.S. Thorne[1]*, A.D. Martin[2], W.J. Stirling[2] and G. Watt[1]

1- Department of Physics and Astronomy, University College London, WC1E 6BT, UK

2- Institute for Particle Physics Phenomenology, University of Durham, DH1 3LE, UK

We consider the impact that can be made on our understanding of parton distributions (PDFs) and QCD from early measurements at the LHCb experiment. The high rapidity values make the experiment uniquely suited to a detailed study of small-x parton distributions and hence will make a significant contribution towards the clarification of both experimental and theoretical uncertainties on PDFs and their applications.

1 Introduction

The kinematic range for particle production at the LHC is shown in Figure 1. The x values of the PDFs are $x_{1,2} = x_0 \exp(\pm y)$, where $x_0 = M/\sqrt{s}$. Smallish $x \sim 0.001 - 0.01$ parton distributions are therefore probed by the standard production processes at the LHC, e.g. W, Z, at central rapidity. However, LHCb, which has a rapidity coverage of $1.8 < y < 4.9$, automatically probes the PDFs at very low (and high) values of x. At the low values the experiment constraints begin to run out and there are also potentially significant theoretical uncertainties.



Figure 1: The kinematic range of x and Q^2 which is probed at the LHC.

We can gain useful information on PDFs and QCD from both total cross-section measurements and ratios of cross-sections at LHCb [1]. The 90% confidence level limit uncertainies using the preliminary MSTW 2007 PDFs [2] are shown in Figure 2. The results are exact but they can be understood more easily using some simplifications. Since $x_1 > x_2$, particularly at high values of y, and since sea quarks die away at high x, we assume $q_1(x_1)\bar{q}_2(x_2) + \bar{q}_1(x_1)q_2(x_2) \approx q_1(x_1)\bar{q}_2(x_2)$. Also at small x_2 we assume that $\bar{u}(x_2) = \bar{d}(x_2)$. We finally assume that the cross-sections are dominated by up and down contributions, which is reasonable for obtaining general results. Using these simplifications we obtain for the cross-section ratios the expressions

$$R_{Z/W} \simeq \frac{A_u\, u(\tilde{x}_1) + A_d\, d(\tilde{x}_1)}{u(x_1) + d(x_1)}, \quad A_\pm \simeq \frac{u_V(x_1) - d_V(x_1)}{u(x_1) + d(x_1)}, \quad R_\pm \simeq \frac{d(x_1)}{u(x_1)},$$

where $A_u = v_u^2 + a_u^2$ and $A_d = v_d^2 + a_d^2$, i.e. the sum of the squares of the vector and axial-vector couplings. The use of $\tilde{x}_1$ illustrates that x values are slightly higher for Z production

---

[*] Royal Society University Research Fellow



Figure 2: The uncertainties on cross-section ratios and total cross-sections at the LHC.

than for W production. In particular the large uncertainty on $R_\pm$ for high y reflects that on the high-x down quark distribution. Similarly for total cross-sections we have

$$\begin{array}{rcl} \sigma_{W^+} & \simeq & u(x_1)\bar{d}(x_2), \quad \sigma_{W^-} \simeq d(x_1)\bar{u}(x_2), \\ \sigma_Z & \simeq & A_u u(x_1)\bar{u}(x_2) + A_d d(x_1)\bar{d}(x_2), \quad \sigma_{\gamma^\star} \simeq 4/9\, u(x_1)\bar{u}(x_2) + 1/9\, d(x_1)\bar{d}(x_2). \end{array} \tag{1}$$

The uncertainty on σ_Z and σ_{W^+} at high y is dominated by the valence up quark and the sea quarks for $x \sim 0.0001$, both of which are limited to a few percent, hence the small uncertainty in Figure 2. The uncertainty on σ_{W^-} is dominated by the down quark for large x_1, as for $A_\pm$. For the virtual photon, $x_{1,2} = (M_\gamma/14{,}000\,\mathrm{GeV})\exp(\pm y)$, so if $M_\gamma \leq 20\,\mathrm{GeV}$, x_2 can be ≤ 0.00001 and x_1 stays away from the valence uncertainty at high x. The increased uncertainty when these extremely small x values are probed is evident in the figure.

2 Potential results from LHCb

Depending on the cross-section under investigation, early data can either reduce the uncertainty on PDFs or test the validity of the order-by-order perturbative expansion. With $1\mathrm{fb}^{-1}$ of data LHCb [3] expects to obtain data for $Z \to \mu^+\mu^-$ for $1.8 < y < 4.9$ with $\sim 1\%$ precision (ignoring luminosity uncertainty) if presented in bins of width 0.1. This is sensitive to the small-x quarks and gluon distribution for $x < 0.001$, but the most precise data will be in the range where HERA data [4] already provide very good constraints. However, if the data are a different shape to the prediction they can highlight problems with fixed order calculations. Illustrated in Figure 3 is pseudo-data shifted by a mulitplicative factor $0.05(y - 3.4)$ compared to the current NLO prediction. The lower (blue) line shows the result of a new fit. It is not possible to obtain good agreement, i.e. $\chi^2 = 103/30$. Hence, this type of result could imply that small-x resummations are important, or that higher twists/saturation is contaminating the current extraction of small-x PDFs from HERA data.

Consider instead the example of 1fb^{-1} data on $\sigma(W^-)/\sigma(W^+)$. (In practice this will be measured as lepton asymmetry, but very similar considerations apply.) The cross-section for $W \to \mu\nu_\mu$ is ten times that for $Z \to \mu^+\mu^-$, but is more difficult to measure with more systematics. We have assumed similar uncertainties, which is perhaps conservative [3]. In this case, assuming the data are consistent with the prediction they immediately become one of the main constraints on the down valence distribution, as shown in Figure 4. Again a discrepancy between data and theory could easily highlight shortcomings, but it is not very obvious what these might be in this case (except perhaps an unexpected large $\bar{u}, \bar{d}$ asymmetry at small x.)



Figure 3: The result of fitting to hypothetical $d\sigma_Z/dy$ data.

Even more dramatic is the constraint that could potentially be made by a measurement of the low-mass Drell-Yan cross-section. In principle the cross-section for e.g. a 14 GeV virtual photon is similar to that for the on-shell Z but the background is far more of a problem. We assume (perhaps slightly optimistically) that for 1fb^{-1} the uncertainty at the lowest y is about twice that for $Z \to \mu^+\mu^-$ but the cross-section (and hence statistics) falls off less quickly with y because the valence quark fall off is not reached. Data of this type for a single mass value produce the extra constraint shown in the right of Figure 4. Since at these values of x the quarks are driven by evolution from the gluon, the constraint is very similar for small-x quarks.



Figure 4: The improved constraint on $d_V(x, Q^2)$ from fitting to $(d\sigma_{W^-}/dy)/(d\sigma_{W^+}/dy)$ data (left) and on $g(x, Q^2)$ from fitting to $(d\sigma_{\gamma^\star}/dy)$ for $M_{\gamma^\star} = 14\,\text{GeV}$ (right).

Since for $x < 0.0001$ the uncertainty on PDFs is already large, a measurement of the low-mass virtual photon cross-section in one mass bin is more inclined to give an improved constraint on PDFs rather than test the theory, unless it is very different from the prediction (more so than required for Z production). However, measurements in the range of mass bins will be an excellent test of small-x evolution. We already have reason to believe that the perturbative series is unstable, where use of the available NNLO matrix elements [5] shows that for $M_{\gamma^\star} < 20\text{GeV}$ there is sizable variation in predictions as one goes from LO to NLO to NNLO, as shown in Figure 5. The instability can be traced to the increasing divergence of terms in the perturbative expansion as one increases the order of the calculation. This has a significant effect on both the small-x PDFs and the matrix elements for the cross-section [6]. Measurements on low-mass Drell-Yan production at LHCb, and also at ATLAS and CMS, will be important for our understanding of the convergence of the perturbative series in QCD. Alternative measurements, e.g. heavy flavour production, could produce complementary information, but are less clean.



Figure 5: The Drell-Yan cross-section as a function of rapidity in various mass bins.

To summarise, suitable measurements at the LHC experiment will place extra constraints on PDFs and QCD which will then feed back into predictions for all quantities. LHCb has a unique reach, and precision measurements at high rapidity can probe some of the most important and interesting regions of parton kinematics, with even limited data from early running making a major impact.

References

[1] Slides:
`http://indico.cern.ch/contributionDisplay.py?contribId=315&sessionId=17&confId=24657`

[2] R. S. Thorne, A. D. Martin, W. J. Stirling and G. Watt, arXiv:0706.0456 [hep-ph].

[3] R. McNulty, these proceedings.

[4] C. Adloff *et al.* [H1 Collaboration], Eur. Phys. J. C **21** (2001) 33 [arXiv:hep-ex/0012053];
S. Chekanov *et al.* [ZEUS Collaboration], Eur. Phys. J. C **21** (2001) 443 [arXiv:hep-ex/0105090].

[5] C. Anastasiou, L. J. Dixon, K. Melnikov and F. Petriello, Phys. Rev. Lett. **91** (2003) 182002 [arXiv:hep-ph/0306192]; Phys. Rev. D **69** (2004) 094008 [arXiv:hep-ph/0312266].

[6] A. D. Martin, W. J. Stirling, R. S. Thorne and G. Watt, in preparation.